

# Habitat Management Guidelines for Bald Eagles in Texas

The following guidelines were developed to help landowners and managers maintain or improve their land for the benefit of the Bald Eagle. Information is also provided so that landowners may recognize and avoid or minimize human-related disturbance to eagles, particularly nesting pairs.

## Nesting Habitat

The protection of an actual nest is important, but so is protection of the nest area and all the surrounding habitat factors that attracted the nesting pair to the area. Once the eagles establish a suitable breeding territory, they will return to the same area year after year, often using several nests within the territory during different years. When a given nest or the tree that it is in falls, a pair generally returns to the same territory to begin another nest. If one member of a pair dies, the nest may go unused for several years and then be recolonized by the surviving member returning with a new mate. Nesting territories can even be inherited by offspring. Therefore, protection of nesting territories should apply to "abandoned" nests for at least five consecutive years of documented nonuse.

The following habitat management guidelines are based on two management zones surrounding each nest site, with certain restrictions recommended for each zone.

### Primary Management Zone For Nest Sites

This zone includes an area extending 750 to 1,500 feet outward in all directions from the nest site. It is recommended that the following activities not occur within this zone:

1. Habitat alteration or change in land use, such as would result from residential, commercial, or industrial development; construction projects; or mining operations.
2. Tree cutting, logging, or removal of trees, either living or dead.

3. Use of chemicals labeled as toxic to fish and wildlife.
4. Placement of above-ground electrical transmission or distribution lines. Since collision with powerlines and electrocution on powerline structures remain an important cause of death, placement of underground lines is recommended near Bald Eagle nests and winter concentration sites.
5. Helicopter or fixed-wing aircraft operation within 500 feet vertical distance or 1,000 feet horizontal distance of the nest site during the nesting season (October-July).
6. Activities which create minimal disturbance, such as hiking, fishing, camping, and bird-watching can be carried out safely during the non-nesting season if there is no physical alteration of the habitat within the zone. Traditional farming, ranching, and hunting activities which are existing practices and have occurred historically on the site can be carried out safely during the non-nesting season as long as habitat alteration is avoided.

Human presence within this zone should be minimized during the nesting season, especially during the early nesting period from October-April. Traditional agricultural activities and low impact recreational activities are generally not a problem even during the nesting season as long as they do not appear to be adversely affecting nesting success, there is no increase in the level of disturbance from historic levels, and physical alteration of the habitat is avoided. However, activities of any kind should be stopped if it becomes apparent that the birds are suffering from disturbance. The key point is whether the activities keep the breeding birds away from the nest, eggs, or

young for extended periods of time. If they do, they are harmful. In general, it is important to protect the nest from human disturbance during very hot or very cold weather, since the parents' absence at these times can be particularly deadly for the eggs or young.

### Secondary Management Zone For Nest Sites

This zone encompasses an area extending outward from the primary zone an additional 750 feet to 1 mile. Recommended restrictions in this zone are intended to protect the integrity of the primary zone and to protect important feeding areas, including the eagle's access to these areas. The following activities are likely to be detrimental to Bald Eagles at any time, and in most cases should be avoided within the secondary zone:

1. Development of new commercial or industrial sites.
2. Construction of multi-story buildings or high-density housing developments between the nest and the eagle's feeding area.
3. Placement of electrical transmission or distribution lines between the nest site and the eagle's feeding area.
4. Construction of new roads, trails, canals, or rights-of-way which would tend to facilitate human access to the eagle nest.
5. Use of chemicals labeled as toxic to wildlife.

Certain activities that involve only minimal alteration or disturbance to the habitat can be carried out safely in the secondary zone during the non-nesting season. Examples of these activities include: minor logging or land clearing, minor construction, seismographic exploration employing explosives, oil well drilling, and low-level aircraft operation. However, these

activities should avoid major alteration or loss of Bald Eagle habitat as much as possible.

If logging is done, it is best to retain as many large trees as possible for roost and perch trees. Retention of at least 10 to 15 live trees per acre is suggested. Ideally, the trees left uncut should be the largest in the stand, preferably those with open crowns and stout lateral limbs. Selective forestry practices such as seedtree, shelterwood, and single tree selection are recommended over clear-cutting.

Minimal disturbance recreational activities (hiking, fishing, camping, picnicking, bird-watching, hunting) and everyday farming and ranching activities that cause no new alteration of habitat can be safely carried out in the secondary zone at any time.

### **Feeding Areas**

The use of toxic chemicals in watersheds and rivers where Bald Eagles feed should be avoided as much as possible. Where agricultural herbicides and pesticides are used within the watershed, label directions should be strictly followed, including those describing proper disposal of rinse water and containers.

Alteration of natural shorelines where Bald Eagles feed should be avoided or minimized as much as possible. Degraded or eroded shorelines should be revegetated whenever possible.

### **Winter Roost Concentration Areas**

Logging or land clearing activity should be avoided within 1,500 feet of a roosting concentration area. Disruptive, noisy, or out-of-the-ordinary land use activities should be avoided near communal roost sites. Normal agricultural activities which have occurred traditionally on the land are generally acceptable near these roost sites as long as they do not appear to be affecting roosting eagles. However, it is best to avoid even normal activities during evening, night, and early morning hours.

### **For More Information**

Landowners and managers can contact the Texas Parks and Wildlife Department, U.S. Fish and Wildlife Service, U.S. Natural Resources Conservation Service (formerly Soil Conservation Service), or Texas Agricultural Extension Service for technical assistance in managing habitat and protecting Bald Eagle nest sites.